

RECOMMENDED VACCINE SCHEDULE FOR CENTRAL COAST OF CALIFORNIA

Disease	Foals & Weanlings	Yearlings	Performance Horses	Pleasure Horses	Broodmares	Comments
WNV (West Nile Virus)	1st dose: 6 mos. 2nd dose: 7 mos.	Yearly, Prior to Mosquito Season	Yearly, Prior to Mosquito Season	Yearly, Prior to Mosquito Season	Yearly, 4-6 weeks before foaling	Vaccinate semi-annually or more frequently (up to every 3 months), depending on risk.
Tetanus	1st dose: 6 mos. 2nd dose: 7 mos.	Yearly	Yearly	Yearly	Yearly, 4-6 weeks before foaling	
3 WAY (EEE, VEE and WEE) Eastern, Venezuelan and Western Equine Encephalomyelitis or Sleeping Sickness	1st dose: 6 mos. 2nd dose: 7 mos.	Yearly, Prior to Mosquito Season	Yearly, Prior to Mosquito Season	Yearly, Prior to Mosquito Season	Yearly, 4-6 weeks before foaling	
Rhino-Flu Equine Influenza and Rhinopneumonitis EHV-1&4 (Intranasal and/or Injectable)	1st dose: 6 mos 2nd dose: 7 mos 3rd dose: 3-4 wks after 2nd dose	Every 3 - 6 mos. Depending on risk of exposure	Every 3 - 6 mos. Depending on risk of exposure	Every 3 - 6 mos. Depending on risk of exposure	At least every 6 mos. with one booster timed 4-6 weeks before foaling	Vaccinate semi-annually or more frequently (up to every 3 months), depending on risk. Recommend alternating Intra-Nasal and Intramuscular routes of administration after initial vaccination series.
Rhinopneumonitis (EHV-1 Abortion Strain - Pneumabort K) <i>* Recommended for broodmares - all others should vaccinate for Rhino-Flu</i>	1st dose: 6 mos 2nd dose: 7 mos 3rd dose: 3-4 wks after 2nd dose	Every 3-4 mos.	Every 3-4 mos.	Every 3-4 mos.	EHV-1 during the 5th, 7th and 9th month of pregnancy (3rd month optional)	Vaccination of mares pre-breeding and 4-6 weeks before foaling with EHV-1 and EHV-4 is recommended
Rabies	1st dose: 6 mos. 2nd dose: 7 mos.	Yearly	Yearly	Yearly	Yearly, 4-6 weeks before foaling	Recommended in areas where wildlife rabies is a problem - Includes the Central Coast
Strangles Strep Equi	1st dose: 6 mos 2nd dose: 7 mos 3rd dose: 3-4 wks after 2nd dose	Every 6 mos.	Optional every 6 mos. if risk is high	Optional every 6 mos. if risk is high	Every 6 mos. with one booster timed 4-6 weeks before foaling	Start initiation of primary series before 6 mos. of age if risk of exposure is high. There are several hot spots for Strangles in SLO County.
Potomac Horse Fever	1st dose: 6 mos. 2nd dose: 7 mos.	Yearly	Yearly	Yearly	Yearly, 4-6 weeks before foaling	No Longer recommended on the West Coast